

AMINO-BENZIMIDAZOLES DERIVATIVES AS INHIBITORS OF RESPIRATORY SYNCYTIAL VIRUS REPLICATION

5 The present invention is concerned with amino-benzimidazole derivatives having
antiviral activity, in particular, having an inhibitory activity on the replication of the
respiratory syncytial virus (RSV). It further concerns their preparation and
compositions comprising them, as well as their use as a medicine.

10 Human RSV or Respiratory Syncytial Virus is a large RNA virus, member of the
family of Paramyxoviridae, subfamily pneumoviridae together with bovine RSV virus.
Human RSV is responsible for a spectrum of respiratory tract diseases in people of all
ages throughout the world. It is the major cause of lower respiratory tract illness during
infancy and childhood. Over half of all infants encounter RSV in their first year of life,
and almost all within their first two years. The infection in young children can cause
15 lung damage that persists for years and may contribute to chronic lung disease in later
life (chronic wheezing, asthma). Older children and adults often suffer from a (bad)
common cold upon RSV infection. In old age, susceptibility again increases, and RSV
has been implicated in a number of outbreaks of pneumonia in the aged resulting in
significant mortality.

20 Infection with a virus from a given subgroup does not protect against a subsequent
infection with an RSV isolate from the same subgroup in the following winter season.
Re-infection with RSV is thus common, despite the existence of only two subtypes, A
and B.


25 Today only three drugs have been approved for use against RSV infection. A first one
is ribavirin, a nucleoside analogue, provides an aerosol treatment for serious RSV
infection in hospitalized children. The aerosol route of administration, the toxicity (risk
of teratogenicity), the cost and the highly variable efficacy limit its use. The other two
30 drugs, RespiGam[®] and palivizumab, polyclonal and monoclonal antibody
immunostimulants, are intended to be used in a preventive way.

Other attempts to develop a safe and effective RSV vaccine have all met with failure
thus far. Inactivated vaccines failed to protect against disease, and in fact in some cases
35 enhanced disease during subsequent infection. Life attenuated vaccines have been tried
with limited success. Clearly there is a need for an efficacious non-toxic and easy to
administer drug against RSV replication.

-2-

Benzimidazoles and imidazopyridines as inhibitors of RSV replication have been described in WO 01/00611, WO 01/00612 and WO 01/00615.

The present invention concerns inhibitors of RSV replication, which can be represented
5 by formula (I):




their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms wherein

Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl substituted with one or more substituents each
10 independently selected from the group consisting of trifluoromethyl, C₃₋₇cycloalkyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkyl-carbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyl-oxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, hydroxy-
15 C₂₋₄-alkyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy, mono- or di(C₁₋₄alkyl)amino-carbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, dioxolanyl optionally substituted with one or two C₁₋₆alkyl radicals, and a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, indolyl, imidazolyl, triazolyl, piperidinyl,
20 homopiperidinyl, piperazinyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with oxo or C₁₋₆alkyl;

G is a direct bond or C₁₋₁₀alkanediyl optionally substituted with one or more substituents individually selected from the group consisting of hydroxy, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹C₁₋₆alkylthio,
25 HO(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-;

R¹ is Ar¹ or a monocyclic or bicyclic heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydro-furanyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl,
30 isoxazolyl, oxadiazolyl, quinolinyl, quinoxalinyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, pyridopyridyl, naphthiridinyl, 1*H*-imidazo[4,5-*b*]pyridinyl, 3*H*-imidazo[4,5-*b*]pyridinyl, imidazo[1,2-*a*]pyridinyl, 2,3-dihydro-1,4-dioxino[2,3-*b*]pyridyl and a radical of formula



- wherein each of said monocyclic or bicyclic heterocycles may optionally be substituted with 1 or where possible more, such as 2, 3, 4 or 5, substituents individually selected from the group of substituents consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)amino, mono-or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino, C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono-or di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n-; each n independently is 1, 2, 3 or 4; one of R^{2a} and R^{3a} is C₁₋₆alkyl and the other one of R^{2a} and R^{3a} is hydrogen; in case R^{2a} is different from hydrogen then R^{2b} is hydrogen or C₁₋₆alkyl, and R^{3b} is hydrogen; in case R^{3a} is different from hydrogen then R^{3b} is hydrogen or C₁₋₆alkyl, and R^{2b} is hydrogen; or R^{2a}, R^{2b}, R^{3a} and R^{3b} all are hydrogen; R^{4a} and R^{4b} can be the same or can be different relative to one another, and are each independently hydrogen or C₁₋₆alkyl; or R^{4a} and R^{4b} taken together may form a bivalent radical of formula -(CH₂)_s-; R⁵ is hydrogen or C₁₋₆alkyl; m is 1 or 2; p is 1 or 2; s is 4 or 5

-4-

Ar¹ is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from halo, hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, and C₁₋₆alkyloxy;

5 Ar² is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from the group consisting of halo, hydroxy, amino, cyano, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkyloxy, amino-sulfonyl, aminocarbonyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, mono- or di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)-aminosulfonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl and C₁₋₄alkoxycarbonyl.

10

The invention relates to the use of a compound of formula (I), or a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex and stereochemically isomeric form thereof, for the manufacture of a medicament for inhibiting RSV replication. Or the invention relates to a method of inhibiting RSV replication in a warm-blooded animal
15 said method comprising the administration of an effective amount of a compound of formula (I), or a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex and stereochemically isomeric form thereof.

15

In a further aspect, this invention relates to novel compounds of formula (I) as well as
20 methods for preparing these compounds.

20

The term 'prodrug' as used throughout this specification and claims means the pharmacologically acceptable derivatives, e.g. esters and amides, such that the resulting biotransformation product of the derivative is the active drug as defined in the
25 compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p. 13-15) describing prodrugs generally, is hereby incorporated. Prodrugs are characterized by a good aqueous solubility and bioavailability, and are readily metabolized into the active inhibitors *in vivo*.

30

The terms 'C₁₋₆alkyl optionally substituted with one or more substituents' such as used in the definition of Q, or 'C₁₋₁₀alkanediyl optionally substituted with one or more substituents' as used in the definition of G are meant to comprise C₁₋₆alkyl radicals respectively C₁₋₁₀alkanediyl radicals having no, one, two or more substituents, for
35 example no, one, two, three, four, five or six substituents, in particular no, one, two or three substituents, further in particular no, one or two substituents. The upper limit of the number of substituents is determined by the number of hydrogen atoms that can be

35

-5-

replaced as well as by the general properties of the substituents such as their bulkiness, these properties allowing the skilled person to determine said upper limit.

As used herein in relation to Q, the term 'wherein each of said heterocycle may
5 optionally be substituted with oxo or C₁₋₆alkyl' is meant to comprise heterocycles substituted with one or more, such up to 3, or up to 2 substituents or with one substituent indepently selected from oxo and C₁₋₆alkyl.

As used in the foregoing and hereinafter, 'polyhaloC₁₋₆alkyl' as a group or part of a
10 group, e.g. in polyhaloC₁₋₆alkyloxy, is defined as mono- or polyhalo substituted C₁₋₆alkyl, in particular C₁₋₆alkyl substituted with up to one, two, three, four, five, six, or more halo atoms, such as methyl or ethyl with one or more fluoro atoms, for example, difluoromethyl, trifluoromethyl, trifluoroethyl. Also included are perfluoro C₁₋₆alkyl
15 groups, which are C₁₋₆alkyl groups whereion all hydrogen atoms are replaced by fluoro atoms, e.g. pentafluoroethyl. In case more than one halogen atom is attached to an alkyl group within the definition of polyhaloC₁₋₄alkyl, the halogen atoms may be the same or different.

Each of the monocyclic or bicyclic heterocycles in the definition of R¹ may optionally
20 be substituted with 1 or where possible more substituents, such as 2, 3, 4 or 5, substituents. In particular, said heterocycles may optionally be substituted with up to 4, up to 3, up to 2 substituents, or up to 1 substituent.

Each Ar¹ or Ar² may be unsubstituted phenyl or phenyl substituted with 1 or more
25 substituents, such as 5 or 4 substituents or, which is preferred, up to 3 substituents, or up to two substituents, or with one substituent.

A hydroxyC₁₋₆alkyl group when substituted on an oxygen atom or a nitrogen atom preferably is a hydroxyC₂₋₆alkyl group wherein the hydroxy group and the oxygen or
30 nitrogen is separated by at least two carbon atoms.

As used herein C₁₋₃alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 3 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl and the like; C₁₋₄alkyl as a group or part of a group defines
35 straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as the group defined for C₁₋₃alkyl and butyl and the like; C₂₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 2 to 4 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl

-6-

and the like; C₁₋₅alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 5 carbon atoms such as the groups defined for C₁₋₄alkyl and pentyl, 1-methylbutyl, 2-methylbutyl, 1-ethylpropyl and the like; C₁₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C₁₋₅alkyl and , hexyl, 2-methylpentyl, 3-methylpentyl and the like; C₁₋₉alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 9 carbon atoms such as the groups defined for C₁₋₆alkyl and heptyl, octyl, nonyl, 2-methylhexyl, 2-methylheptyl and the like; C₁₋₁₀alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 10 carbon atoms such as the groups defined for C₁₋₉alkyl and decyl, 2-methylnonyl and the like.

C₃₋₇cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

C₂₋₅alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 2 to 5 carbon atoms such as, for example, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,2-propanediyl, 2,3-butanediyl, 1,5-pentanediyl and the like, C₁₋₄alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like; C₁₋₆alkanediyl is meant to include C₁₋₄alkanediyl and the higher homologues thereof having from 5 to 6 carbon atoms such as, for example, 1,5-pentanediyl, 1,6-hexanediyl and the like; C₁₋₁₀alkanediyl is meant to include C₁₋₆alkanediyl and the higher homologues thereof having from 7 to 10 carbon atoms such as, for example, 1,7-heptanediyl, 1,8-octanediyl, 1,9-nonanediyl, 1,10-decanediyl and the like.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom. The term (=N-OH) forms a hydroxylimine moiety when attached to a carbon atom.

The term halo is generic to fluoro, chloro, bromo and iodo.

It should be noted that the radical positions on any molecular moiety used in the definitions may be anywhere on such moiety as long as it is chemically stable.

-7-

Radicals used in the definitions of the variables include all possible isomers unless otherwise indicated. For instance pyridyl includes 2-pyridyl, 3-pyridyl and 4-pyridyl; pentyl includes 1-pentyl, 2-pentyl and 3-pentyl.

- 5 When any variable occurs more than one time in any constituent, each definition is independent.

Whenever used hereinafter, the term 'compounds of formula (I)', or 'the present compounds' or similar term is meant to include the compounds of general formula (I),
10 their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms. An interesting subgroup of the compounds of formula (I) or any subgroup thereof are the *N*-oxides, salts and all the stereoisomeric forms of the compounds of formula (I).

- 15 It will be appreciated that some of the compounds of formula (I) may contain one or more centers of chirality and exist as stereochemically isomeric forms.

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible compounds made up of the same atoms bonded by the same sequence of bonds
20 but having different three-dimensional structures which are not interchangeable, which the compounds of formula (I) may possess.

Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms which said
25 compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in pure form or in admixture with each other are intended to be embraced within the scope of the present invention.

30 Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i. e. minimum 90% of one isomer and maximum
35 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and

'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric excess, respectively the diastereomeric excess of the mixture in question.

5 Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyl-tartaric acid, ditoluoyltartaric acid and camphosulfonic acid. Alternatively, enantiomers
10 may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These
15 methods will advantageously employ enantiomerically pure starting materials.

The diastereomeric racemates of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column
20 chromatography.

For some of the compounds of formula (I), their prodrugs, N-oxides, salts, solvates, quaternary amines, or metal complexes and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined.
25 A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction.

The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but
30 different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases, which are
35 non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

-9-

The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic (i.e. hydroxybutanedioic acid), tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-*D*-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkyl halide, aryl halide or arylalkyl halide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen.

Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

-10-


The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

It will be appreciated that the compounds of formula (I) may have metal binding,
5 chelating, complex forming properties and therefore may exist as metal complexes or metal chelates. Such metalated derivatives of the compounds of formula (I) are intended to be included within the scope of the present invention.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such
10 forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.


Any subgroup of compounds of formula (I) specified herein is meant to also comprise the prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and
15 stereochemically isomeric forms of this subgroup of compounds of formula (I).

One embodiment of the present invention concerns compounds of formula (I-a):



wherein Q, R⁵, G and R¹ are as specified above or as in any of the subgroups of
20 compounds specified herein; and
R^{2a} is C₁₋₆alkyl;
R^{2b} is hydrogen or C₁₋₆alkyl.

Another embodiment of the present invention concerns compounds of formula (I-b):




wherein Q, R⁵, G and R¹ are as specified above or as in any of the subgroups of
25 compounds specified herein; and
R^{3a} is C₁₋₆alkyl;
R^{3b} is hydrogen or C₁₋₆alkyl.

30

-11-

Another embodiment of the present invention concerns compounds of formula (I-c):



wherein Q, R⁵, G and R¹ are as specified above or as in any of the subgroups of compounds specified herein.

It is to be understood that the above defined subgroups of compounds of formulae (I-a), (I-b), etc. as well as any other subgroup defined herein, are meant to also comprise any prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms of such compounds.

Particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein G is C₁₋₁₀alkanediyl, more in particular wherein G is methylene.

Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein (a) R¹ is other than Ar¹; or wherein (b) R¹ is Ar¹ or a monocyclic heterocycle, which is as specified in the definitions of the compounds of formula (I) or any of the subgroups thereof.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein


- (c) R¹ is pyridyl optionally substituted with 1 or 2 substituents independently selected from the group consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)amino, mono-or di(C₁₋₆alkyl)amino-C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino, C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono-or di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n-; or more in particular
- (d) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy, C₁₋₆alkyl, halo, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy and (C₁₋₆alkyloxy)C₁₋₆alkyloxy; preferably wherein
- (e) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy, C₁₋₆alkyl, halo and C₁₋₆alkyloxy; or wherein

-12-

- (f) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy and C₁₋₆alkyl; more preferably wherein
- (g) R¹ is pyridyl substituted with hydroxy and C₁₋₆alkyl; or more preferably wherein
- (h) R¹ is pyridyl substituted with hydroxy and methyl; or wherein
- 5 (i) R¹ is 3-hydroxy-6-methylpyrid-2-yl.

Further embodiments comprise those compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein

- (j) R¹ is Ar¹, quinolinyl, benzimidazolyl, a radical of formula



10

pyrazinyl, or pyridyl; or wherein

- (k) R¹ is Ar¹, quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl, or pyridyl;
- 15 wherein each of the radicals in (j) and (k) may optionally be substituted with the substituents specified in the definition of the compounds of formula (I) and in particular pyridyl may be substituted as specified above in (a) to (i).
- Further embodiments comprise those compounds of formula (I) or any of the subgroups
- 20 of compounds of formula (I) wherein
- (l) R¹ is Ar¹, quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl, or pyridyl, wherein each of these radicals may optionally be substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy, (C₁₋₆alkyloxy)C₁₋₆alkyloxy; or more
- 25 specifically wherein
- (m) R¹ is Ar¹, quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl, or pyridyl, wherein each of these radicals may optionally be substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, benzyloxy; or more specifically wherein
- 30 (n) R¹ is phenyl optionally substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy; quinolinyl; a radical (c-4) wherein m is 2, optionally substituted with up to two radicals selected from C₁₋₆alkyl; benzimidazolyl optionally substituted with C₁₋₆alkyl; pyridyl optionally substituted with one or two radicals selected from hydroxy, halo, C₁₋₆alkyl,
- 35 benzyloxy and C₁₋₆alkyloxy, pyrazinyl optionally substituted with up to three

-13-

radicals selected from C₁₋₆alkyl; or pyridyl substituted or optionally substituted as specified above in (a) – (i); or wherein

- (o) R¹ is phenyl optionally substituted with one or two radicals selected from the group consisting of halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy; or
- 5 (p) R¹ is quinoliny; or
- (q) R¹ is a radical (c-4) wherein m is 2, optionally substituted with up to two radicals selected from C₁₋₆alkyl; or
- (r) R¹ is benzimidazolyl optionally substituted with C₁₋₆alkyl; pyridyl optionally substituted with one or two radicals selected from hydroxy, halo, C₁₋₆alkyl,
- 10 benzyloxy and C₁₋₆alkyloxy; or
- (s) R¹ is pyrazinyl optionally substituted with up to three radicals selected from C₁₋₆alkyl.

Preferred subgroups of compounds of formula (I) or any of the subgroups of
15 compounds of formula (I) are those wherein G is a direct bond or methylene and R¹ is as specified above in (a) – (s). Further preferred are the compounds of formula (I) or any of the subgroups specified herein wherein G is a direct bond and R¹ is a radical (c-4), in particular wherein m is 2, optionally substituted with up to two radicals selected from C₁₋₆alkyl. Further preferred are the compounds of formula (I) or any of
20 the subgroups specified herein wherein G is methylene and R¹ is as specified above in (a) – (s), but is other than a radical (c-4).

Other embodiments comprise those compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein, wherein R⁵ is hydrogen.
25

Other embodiments comprise those compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein, wherein:

- (a) Q is Ar², C₃₋₇cycloalkyl or C₁₋₆alkyl substituted with one or two substituents each independently selected from the group consisting of substituents mentioned in the
30 definition of the compounds of formula (I) or of any subgroup thereof; or in particular
- (b) Q is Ar², C₃₋₇cycloalkyl or C₁₋₆alkyl substituted with one substituent selected from the group consisting of substituents mentioned in the definition of the compounds of formula (I) or of any subgroup thereof, and said C₁₋₆alkyl is optionally further
35 substituted with one hydroxy; or
- (c) Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl optionally substituted with one or two substituents each independently selected from the group consisting of trifluoromethyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²(CH₂)_noxy, hydroxy-

-14-

- carbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyloxy, hydroxy-C₂₋₄-alkyloxy, mono- or di(C₁₋₄alkyl)amino-carbonyl, dioxolanyl optionally substituted with one or two C₁₋₆alkyl radicals, and a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, indolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with up to two substituents independently selected from oxo and C₁₋₆alkyl; or
- (d) Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl optionally substituted with one substituent selected from trifluoromethyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylcarbonyloxy, hydroxy-C₂₋₄-alkyloxy, mono- or di(C₁₋₄alkyl)-aminocarbonyl, dioxolanyl optionally substituted with one or two C₁₋₆alkyl radicals, and a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, indolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with up to two substituents independently selected from oxo and C₁₋₆alkyl, and said C₁₋₆alkyl is optionally further substituted with one hydroxy; or
- (e) Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl optionally substituted with one or two substituents each independently selected from the group consisting of Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, aminocarbonyl, C₁₋₄alkoxycarbonyl, hydroxy-C₂₋₄-alkyloxy, dioxolanyl substituted with two C₁₋₆alkyl radicals, and a heterocycle selected from the group consisting of pyrrolidinyl, indolyl, imidazolyl, piperidinyl, piperazinyl, and pyridyl, wherein each of said heterocycle may optionally be substituted with up to two substituents independently selected from oxo and C₁₋₆alkyl; or
- (f) Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl optionally substituted with Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, aminocarbonyl, C₁₋₄alkoxycarbonyl, hydroxy-C₂₋₄-alkyloxy, dioxolanyl substituted with two C₁₋₆alkyl radicals, or a heterocycle selected from pyrrolidinyl, indolyl, imidazolyl, piperidinyl, piperazinyl, and pyridyl, wherein each of said heterocycle may optionally be substituted with up to two substituents independently selected from oxo and C₁₋₆alkyl, and said C₁₋₆alkyl is optionally further substituted with one hydroxy; or
- (g) Q is Ar², C₃₋₇cycloalkyl, or C₁₋₆alkyl optionally substituted with Ar², with one or two hydroxyl groups, with C₁₋₄alkoxy, C₁₋₄alkylthio, aminocarbonyl, C₁₋₄alkoxycarbonyl, hydroxy-C₂₋₄-alkyloxy, dioxolanyl substituted with two C₁₋₆alkyl

-15-

- radicals, or a heterocycle selected from pyrrolidinyl, indolyl, imidazolyl, piperidinyl, piperazinyl, and pyridyl, wherein each of said heterocycle may optionally be substituted with two substituents independently selected from oxo and C₁₋₆alkyl; or
- 5 (h) Q is C₁₋₆alkyl optionally substituted with Ar², with one or two hydroxyl groups, with C₁₋₄alkoxy, C₁₋₄alkylthio, aminocarbonyl, C₁₋₄alkoxycarbonyl, or a heterocycle selected from pyrrolidinyl, imidazolyl, piperidinyl and piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or C₁₋₆alkyl, or with oxo and C₁₋₆alkyl; or
- 10 (i) Q is Ar².

Interesting subgroups among the subgroups mentioned in the previous paragraph are those wherein Ar² is phenyl or phenyl substituted with 1, 2 or 3 substituents or with 1 or 2 substituents, or preferably with one substituent selected from halo, hydroxy, amino, cyano, hydroxyc₁₋₆alkyl, aminoc₁₋₆alkyl, C₁₋₆alkyloxy and aminosulfonyl.

15 Further interesting subgroups among the subgroups mentioned in the previous paragraph are those wherein Ar² is phenyl or phenyl substituted with 1, 2 or 3 substituents or with 1 or 2 substituents, or preferably with one substituent selected from amino, cyano, hydroxyc₁₋₆alkyl, aminoc₁₋₆alkyl and aminosulfonyl.

20 In particular, Ar¹ is phenyl or phenyl substituted with 1, 2, 3 substituents or with 1, 2 substituents selected from those mentioned in the definition of the compounds of formula (I) or of any subgroup thereof.

25 Ar² is phenyl or phenyl substituted with 1, 2, 3 substituents or with 1, 2 substituents selected from the group consisting of those mentioned in the definition of the compounds of formula (I) or of any subgroup thereof.

In the group of compounds of formula (I) or in any of the subgroups of compounds of formula (I):

30

- (a) Ar¹ preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo, hydroxy, C₁₋₆alkyl, hydroxyc₁₋₆alkyl, trifluormethyl, and C₁₋₆alkyloxy;
- (b) Ar¹ more preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo, hydroxy, C₁₋₆alkyl and C₁₋₆alkyloxy;
- 35

-16-


(c) Ar¹ more preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo and C₁₋₆alkyl.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Ar² is as defined for Ar¹.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein one of R^{2a} and R^{3a} is C₁₋₆alkyl and the other one of R^{2a} and R^{3a} is hydrogen; in case R^{2a} is different from hydrogen then R^{2b} is C₁₋₆alkyl, and R^{3b} is hydrogen; in case R^{3a} is different from hydrogen then R^{3b} is C₁₋₆alkyl, and R^{2b} is hydrogen.

Preferred compounds are those compounds listed in tables 1 through 3, more in particular the compound numbers 1 to 11 and 25 to 28.

The compounds of formula (I) or any of the subgroups thereof can be prepared as in the following reaction schemes.



In these schemes Q, G, R¹, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R⁵ have the meanings defined above for the compounds of formula (I) or of any of the subgroups thereof. W is an appropriate leaving group, preferably it is chloro or bromo. The reactions of these schemes can be typically conducted in a suitable solvent such as an ether, e.g. THF, a halogenated hydrocarbon, e.g. dichloromethane, CHCl₃, toluene, a polar aprotic solvent such as DMF, DMSO, DMA and the like. A base may be added to pick up the acid that is

-17-

liberated during the reaction. If desired, certain catalysts such as iodide salts (e.g. KI) may be added.

Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions, comprising those described hereinafter.


Compounds of formula (I) wherein R⁵ is hydrogen may be converted to corresponding compounds of formula (I) wherein is other than hydrogen by an N-alkylation reaction which may be conducted under similar conditions as described above for the conversion of (II) or (IV) to (I).

Compounds wherein Q is C₁₋₆alkyl substituted with C₁₋₄alkoxycarbonyl can be reduced with e.g. LiAlH₄ to the corresponding compounds wherein Q is C₁₋₆alkyl substituted with hydroxy. The same starting materials can be reacted with an amine to the corresponding amides.


Compounds of formula (I) wherein Q is Ar having a cyano or a cyanoC₁₋₅alkyl substituent can be reduced with hydrogen in the presence of a suitable catalyst such as Raney nickel, to yield the corresponding methyleneamine or aminoC₁₋₆alkyl substituents.

A number of the intermediates used to prepare the compounds of formula (I) are known compounds or are analogs of known compounds, which can be prepared following modifications of art-known methodologies readily accessible to the skilled person. A number of preparations of intermediates are given hereafter in somewhat more detail.


The intermediates of formula (II) and (IV) can be prepared as outlined in the following reaction schemes.



-18-



5



- In a first step, a diaminobenzene (VI) is cyclized with urea, preferably in a suitable solvent, e.g. xylene, to yield a benzimidazolone (VII). The latter is converted to a benzimidazole derivative (VIII) wherein W is a leaving group as specified above, in particular by reaction of (VII) with a suitable halogenating agent, for example POCl₃. The resulting intermediate (VIII) is reacted with an amine derivative (IX) in an N-alkylation reaction to obtain an intermediate (II).
- Intermediate (VIII) can be similarly reacted with an amine (XII) to yield intermediate (IV). The above-mentioned reactions are conducted in a suitable solvent and, if desired, in the presence of a base.

- The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxy-peroxy acid or halo substituted benzenecarboxy-peroxy acid, e.g. 3-chlorobenzenecarboxy-peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g.

-19-

t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

- 5 Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.
- 10 The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.
- 20
- 25 In a further aspect, the present invention concerns a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I) as specified herein, or a compound of any of the subgroups of compounds of formula (I) as specified herein, and a pharmaceutically acceptable carrier. A therapeutically effective amount in this context is an amount sufficient to prophylactically act against, to stabilize or to reduce viral infection, and in particular RSV viral infection, in infected subjects or subjects being at risk of being infected. In still a further aspect, this invention relates to a process of preparing a pharmaceutical composition as specified herein, which comprises intimately mixing a pharmaceutically acceptable carrier with a therapeutically effective amount of a compound of formula (I), as specified herein, or of a compound of any of the subgroups of compounds of formula (I) as specified herein.
- 30
- 35

-20-

Therefore, the compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form or metal complex, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

The compounds of the present invention may also be administered via oral inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder, a solution being preferred. Any system developed for the delivery of solutions, suspensions or dry powders via oral inhalation or insufflation are suitable for the administration of the present compounds.

-21-

Thus, the present invention also provides a pharmaceutical composition adapted for administration by inhalation or insufflation through the mouth comprising a compound of formula (I) and a pharmaceutically acceptable carrier. Preferably, the compounds of the present invention are administered via inhalation of a solution in nebulized or aerosolized doses.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, injectable solutions or suspensions and the like, and segregated multiples thereof.

The compounds of formula (I) show antiviral properties. Viral infections treatable using the compounds and methods of the present invention include those infections brought on by ortho- and paramyxoviruses and in particular by human and bovine respiratory syncytial virus (RSV). A number of the compounds of this invention moreover are active against mutated strains of RSV. Additionally, many of the compounds of this invention show a favorable pharmacokinetic profile and have attractive properties in terms of bioavailability, including an acceptable half-life, AUC and peak values and lacking unfavourable phenomena such as insufficient quick onset and tissue retention.

The *in vitro* antiviral activity against RSV of the present compounds was tested in a test as described in the experimental part of the description, and may also be demonstrated in a virus yield reduction assay. The *in vivo* antiviral activity against RSV of the present compounds may be demonstrated in a test model using cotton rats as described in Wyde et al. (Antiviral Research (1998), 38, 31-42).

Due to their antiviral properties, particularly their anti-RSV properties, the compounds of formula (I) or any subgroup thereof, their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms, are useful in the treatment of individuals experiencing a viral infection, particularly a RSV infection, and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses, in particular the respiratory syncytial virus.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines. Said use as a medicine or method of treatment comprises the systemic administration to viral infected subjects or to subjects susceptible to viral infections of
5 an amount effective to combat the conditions associated with the viral infection, in particular the RSV infection.

The present invention also relates to the use of the present compounds or any subgroup thereof in the manufacture of a medicament for the treatment or the prevention of viral
10 infections, particularly RSV infection.

The present invention furthermore relates to a method of treating a warm-blooded animal infected by a virus, or being at risk of infection by a virus, in particular by RSV, said method comprising the administration of an anti-virally effective amount of a
15 compound of formula (I), as specified herein, or of a compound of any of the subgroups of compounds of formula (I), as specified herein.

In general it is contemplated that an antiviral effective daily amount would be from 0.01 mg/kg to 500 mg/kg body weight, more preferably from 0.1 mg/kg to 50 mg/kg
20 body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective
25 daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.

Also, the combination of another antiviral agent and a compound of formula (I) can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I), and (b) another antiviral compound, as a combined
35 preparation for simultaneous, separate or sequential use in antiviral treatment. The different drugs may be combined in a single preparation together with pharmaceutically

-23-

acceptable carriers. For instance, the compounds of the present invention may be combined with interferon-beta or tumor necrosis factor-alpha in order to treat or prevent RSV infections.

5 Examples

The following examples are intended to illustrate the present invention and not to limit it thereto. The terms 'compound 1, compound 4, etc. used in these examples refers to the same compounds in the tables.

10 The compounds were identified by LC/MS using the following equipment:


15 LCT: electrospray ionisation in positive mode, scanning mode from 100 to 900 amu; Xterra MS C18 (Waters, Milford, MA) 5 µm, 3.9 x 150 mm); flow rate 1 ml/min. Two mobile phases (mobile phase A: 85% 6.5mM ammonium acetate + 15% acetonitrile; mobile phase B: 20% 6.5 mM ammonium acetate + 80% acetonitrile) were employed to run a gradient from 100 % A for 3 min to 100% B in 5 min., 100% B for 6 min to 100 % A in 3 min, and equilibrate again with 100 % A for 3 min).

20 ZQ: electrospray ionisation in both positive and negative (pulsed) mode scanning from 100 to 1000 amu; Xterra RP C18 (Waters, Milford, MA) 5 µm, 3.9 x 150 mm); flow rate 1 ml/min. Two mobile phases (mobile phase A: 85% 6.5mM ammonium acetate + 15% acetonitrile; mobile phase B: 20% 6.5 mM ammonium acetate + 80% acetonitrile) were employed to run a gradient condition from 100 % A for 3 min to 100% B in 5 min., 100% B for 6 min to 100 % A in 3 min, and equilibrate again with 100 % A for 3 min).
25 min).

Example 1: Preparation of dimethylbenzimidazolamines

-24-

Scheme A



Preparation of intermediate a-2:

SOCl₂ (14 ml) was added drop wise to a solution of (3-benzyloxy-6-methyl-pyridin-2-yl)-methanol (0.0606 mol) at 5°C. The reaction was stirred at room temperature for 3 hours. The solvent was evaporated under reduced pressure. The residue was taken up in diethyl ether. The precipitate was filtered off and dried, yielding 16.9g of **a-2** (98%, melting point: 182°C).

Preparation of intermediate a-4:

A mixture of 2-chloro-4,6-dimethyl-1H-benzimidazole (0.083 mol), **a-2** (0.0913 mol) and K₂CO₃ (0.332 mol) in dimethylformamide (100ml) was stirred at room temperature for 24 hours. H₂O was then added. The mixture was extracted three times with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated at 30°C under reduced pressure. The residue was taken up in CH₃CN/diisopropylether. The precipitate was filtered off and dried, yielding 16.8g of **a-4** (52%, melting point: 155°C).

Preparation of intermediate a-5:

A mixture of **a-4** (0.0007 mol) and 3-piperidin-1-yl-propylamine (0.003 mol) was stirred at 130°C for 2 hours. The residue was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding: 0.174g of [1-(3-benzyloxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzimidazol-2-yl]-(3-piperidin-1-yl-propyl)-amine (46%).


Preparation of final compound a-6:

-25-

A mixture of [1-(3-benzyloxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-yl]-(3-piperidin-1-yl-propyl)-amine (0.0003 mol) and Pd/C (0.06g) in CH₃OH (10ml) was hydrogenated at room temperature for 1 hour under a 3 bar pressure, then filtered over celite. The filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 89/10/1; 10μm). The pure fractions were collected and the solvent was evaporated. The residue (0.084g) was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding 0.073g of 2-[4,6-dimethyl-2-(3-piperidin-1-yl-propylamino)-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (Compound 1, 51%, melting point: > 260°C).

Example 2: Preparation of dihydroxyalkyl substituted dimethylbenzimidazoleamines

Scheme B



*Preparation of intermediate **b-3**:*

A mixture of **b-1** (0.0014 mol) and **b-2** (0.0012 mol) was stirred at 130°C for 3 hours, then stirred at 160°C for 2 hours, cooled down to room temperature and taken up in CH₂Cl₂. The organic layer was washed with a 10% solution of K₂CO₃, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 97/3/0.1). The pure fractions were collected and the solvent was evaporated, yielding 0.55g of intermediate **b-3** (81%).

*Preparation of compound **b-4**:*

A mixture of **b-3** (0.0011 mol) and Pd/C (0.18g) in CH₃OH (10ml) was hydrogenated for 1 hour under a 3 bar pressure, then filtered over celite. Celite was rinsed with

-26-


CH₃OH. The filtrate was concentrated under reduced pressure. The residue (0.47g) was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding 0.27g of 2-{2-[(2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-amino]-4,6-dimethyl-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol (compound 21, 60%, melting point: 225°C).

5 *Preparation of final compound b-5:*

A mixture of 2-{2-[(2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-amino]-4,6-dimethyl-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol (0.0005 mol) in a 3N solution of HCl (15ml) and tetrahydrofuran (15ml) was stirred for 4 hours. Tetrahydrofuran was evaporated under reduced pressure. The mixture was basified with K₂CO₃ (powder).

10 The aqueous layer was saturated with K₂CO₃ (powder). A solution of CH₂Cl₂/CH₃OH (90/10) was added. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated. The residue (0.17g, 88%) was crystallized from CH₃CN/diisopropylether. The precipitate was filtered off and dried. Yielding: 0.085g of
15 3-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-propane-1,2-diol (compound 4, melting point: 205°C).

Example 3: Preparation of hydroxyalkyl substituted dimethylbenzimidazoleamines



20 *Preparation of intermediate c-3:*

A mixture of **c-2** (0.004 mol) and **c-1** (0.006 mol) was stirred at 130°C for 12 hours, and then taken up in CH₂Cl₂. The organic layer was washed with a 10% solution of K₂CO₃, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue (0.6g) was purified by column chromatography over silica gel (eluent:

25 CH₂Cl₂/CH₃OH/NH₄OH 95/5/0.2; 10μm). The pure fractions were collected and the solvent was evaporated, yielding 0.4g of intermediate **c-3** (38%).

Preparation of compound c-5:

-27-


A mixture of **c-3** (0.0015 mol), **c-4** (0.0016 mol) and K_2CO_3 (0.0052 mol) in dimethyl-formamide (20ml) was stirred at 70°C for 4 hours. The solvent was evaporated until dryness. The residue was taken up in CH_2Cl_2 . The organic layer was washed with H_2O , dried (over $MgSO_4$), filtered and the solvent was evaporated. The residue (0.81g) was purified by column chromatography over silica gel (eluent: toluene/2-propanol/ NH_4OH 90/10/0.5; 10 μ m). The pure fractions were collected and the solvent was evaporated. The residue (0.12g) was crystallized from 2-propanol/ CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 0.12g of 3-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-propionic acid ethyl ester (compound 12, 21%, melting point: 180°C).

Preparation of final compound c-6:

$LiAlH_4$ (0.0003 mol) was added portion wise at 5°C to a mixture of 3-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-propionic acid ethyl ester (0.0001 mol) in tetrahydrofuran (10ml) under N_2 flow. The mixture was stirred at 5°C for 1 hour, then at room temperature for 3 hours. Ethylacetate and H_2O were added. The mixture was extracted with ethylacetate. The organic layer was separated, dried (over $MgSO_4$), filtered and the solvent was evaporated until dryness. The residue was crystallized from 2-propanone/ CH_3CN / diisopropylether. The precipitate was filtered off and dried, yielding 0.025g of 2-[2-(3-hydroxypropylamino)-4,6-dimethyl-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (compound 7, 73%, melting point: 170°C).

Example 4: Preparation of amidoalkyl substituted dimethylbenzimidazoleamines

Scheme D



A mixture of 3-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-propionic acid ethyl ester (0.0001 mol) in a saturated solution of NH_3 in CH_3OH (10ml) was stirred at 70°C for 6 hours. The solvent was evaporated until dryness. The residue (0.05g) was purified by column chromatography over silica gel (eluent: $CH_2Cl_2/CH_3OH/NH_4OH$ 88/12/1; 10 μ m). The pure fractions were collected and the solvent was evaporated. The residue (0.022g, 48%) was


-28-

crystallized from 2-propanone/CH₃CN/diisopropylether. The precipitate was filtered off and dried, yielding 0.014g of 3-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-propionamide **d-2** (compound 8, 30%, melting point: 229°C).

5

Example 5: Preparation of aryl substituted dimethylbenzimidazoleamines

Scheme E



Preparation of intermediate e-3:

- 10 A mixture of **e-1** (0.0022 mol) and **e-2** (0.0023 mol) was stirred at 130°C for 1 hour, then cooled down to room temperature and taken up in CH₂Cl₂. The precipitate was filtered. The mother layer was evaporated. The residue (0.522g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 99/1/0.1 to 90/10/1; 5μm). The pure fractions were collected and the solvent was evaporated,
- 15 yielding 0.36g of intermediate **e-3** (62%).

Preparation of compound e-5:

4-[1-(3-Hydroxy-6-methyl-pyridin-2-ylmethyl)-4,6-dimethyl-1H-benzoimidazol-2-ylamino]-benzonitrile (compound 20, melting point: > 260°C) was prepared analogous to the procedure described for **e-5**.

- 20 *Preparation of final compound e-6:*

- Raney Nickel (0.2g) was added to a mixture of **e-5** (0.0001 mol) in a saturated solution of NH₃ in CH₃OH (20ml). The mixture was hydrogenated at room temperature for 3 hours under a 5 bar pressure, then filtered over celite. Celite was rinsed with H₂O. The filtrate was evaporated until dryness. The residue (0.07g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 85/15/1; 10μm). The pure fractions were collected and the solvent was evaporated. The residue (0.042g, 84%) was crystallized from 2-propanone/CH₃CN/diisopropylether. The precipitate was
- 25 filtered off and dried, yielding 0.022g of 2-[2-(4-aminomethyl-phenylamino)-4,6-


-29-

dimethyl-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol, **e-6** (compound 9, 44%, melting point: 255°C).

Example 6: Preparation of aryl substituted dimethylbenzimidazoleamines

5

Scheme F



Preparation of intermediate f-3:

The intermediate **f-3** was prepared analogous to the procedure described for intermediate **e-3**.
10

Preparation of intermediates f-5 and f-6:

The mixture of intermediates **f-5** and **f-6** has been prepared analogous to the procedure described for **c-5**.

Preparation of f-7 and f-8:


15 The compounds **f-7** and **f-8** have been prepared analogous to the procedure described for **e-6**, yielding 0.18g of fraction 1 (10%) and 0.36g of fraction 2 (20%). Fraction 1 was transformed in acetate and crystallized from 2-propanone/CH₃CN/diisopropylether. The precipitate was filtered off and dried, yielding 0.013g of **f-8** (7.5%, 1 CH₃CO₂H, melting point: 171°C). Fraction 2 was dissolved in
20 2-propanol/HCl and converted into the hydrochloric acid salt. The precipitate was filtered off and dried. The residue was crystallized from 2-propanol/diisopropylether.

-30-

The precipitate was filtered off and dried, yielding 0.021g of **f-7** (compound 26, 10.4%, 4 HCl, melting point: 213°C).

5 The following tables list compounds of the present invention that were prepared analogous to any one of the above mentioned synthesis schemes.

Table 1



Comp. No.	R	Activity	Mass Spectroscopy	Melting point	Synthesis Scheme / salt form
1		7.7	MH ⁺ = 408	> 260°C	A
2		7.7	MH ⁺ = 394	225°C	A
3		7.6	MH ⁺ = 423	225°C	A
4		7.4	MH ⁺ = 357	205°C	B
5		7.4	MH ⁺ = 394	185°C	A
6		7.2	MH ⁺ = 394	230°C	A
7		6.8	MH ⁺ = 341	170°C	C
8		6.8	MH ⁺ = 354	229°C	D
9		6.6	MH ⁺ = 388	255°C	E
10		6.3	MH ⁺ = 391	255°C	A

-31-














Comp. No.	R	Activity	Mass Spectroscopy	Melting point	Synthesis Scheme / salt form
11		6.2	$MH^+ = 377$	$> 260^\circ C$	A
12		5.5	$MH^+ = 383$	$180^\circ C$	C
13		5.5	$MH^+ = 402$	$171^\circ C$	F / HCl
14		4.0	$MH^+ = 374$	$254^\circ C$	E
15		5.8	$MH^+ = 374$	$208^\circ C$	C
16		4.0	$MH^+ = 355$	$206^\circ C$	C
17		4.0	$MH^+ = 365$	$176^\circ C$	F
18		4.0	$MH^+ = 387$	$232^\circ C$	C
19		4.0	$MH^+ = 384$	$> 260^\circ C$	E
20		4.0	$MH^+ = 397$	$225^\circ C$	B
21		4.0	$MH^+ = 408$	$170^\circ C$	A
22		4.2	$MH^+ = 466$	$> 260^\circ C$	A

Table 2



-32-















Comp. No.	R	Activity	Mass Spectroscopy	Melting point	Synthesis scheme / salt form
23		5.2	$MH^+ = 402$	171°C	F / acetate
24		4.0	$MH^+ = 379$	222°C	F

Table 3



Comp. No.	R	Activity	Mass Spectroscopy	Melting point	Synthesis scheme
25		6.5	$MH^+ = 380$	195°C	C
26		6.5	$MH^+ = 395$	190°C	C
27		6.3	$MH^+ = 329$	170°C	C
28		6.2	$MH^+ = 366$	190°C	C
29		5.6	$MH^+ = 389$	215°C	C
30		5.4	$MH^+ = 369$	221°C	C
31		5.3	$MH^+ = 341$	182°C	C
32		5.0	$MH^+ = 313$	210°C	C
33		5.0	$MH^+ = 355$	185°C	C
34		5.0	$MH^+ = 419$	180°C	C
35		4.9	$MH^+ = 380$	175°C	C

-33-

Comp. No.	R	Activity	Mass Spectroscopy	Melting point	Synthesis scheme
36		4.8	MH ⁺ = 343	205°C	C
37		4.0	MH ⁺ = 369	215°C	C
38		4.3	MH ⁺ = 340	220°C	C
39		4.0	MH ⁺ = 398	245°C	C
40		4.3	MH ⁺ = 360	225°C	C
41		4.0	MH ⁺ = 377	245°C	C
42		4.0	MH ⁺ = 377	250°C	C
43		4.1	MH ⁺ = 343	215°C	C

Example 7: *In vitro* screening for activity against Respiratory Syncytial Virus.

The percent protection against cytopathology caused by viruses (antiviral activity or EC₅₀) achieved by tested compounds and their cytotoxicity (CC₅₀) are both calculated from dose-response curves. The selectivity of the antiviral effect is represented by the selectivity index (SI), calculated by dividing the CC₅₀ (cytotoxic dose for 50% of the cells) by the EC₅₀ (antiviral activity for 50 % of the cells). The tables in the above experimental part list the category to which each of the prepared compounds belong : Compounds belonging to activity category “A” have an pEC₅₀ (-log of EC₅₀ when expressed in molar units) equal to or more than 6. Compounds belonging to activity category “B” have a pEC₅₀ value below 6.

Automated tetrazolium-based colorimetric assays were used for determination of EC₅₀ and CC₅₀ of test compounds. Flat-bottom, 96-well plastic microtiter trays were filled with 180 µl of Eagle's Basal Medium, supplemented with 5 % FCS (0% for FLU) and 20 mM Hepes buffer. Subsequently, stock solutions (7.8 x final test concentration) of compounds were added in 45 µl volumes to a series of triplicate wells so as to allow simultaneous evaluation of their effects on virus- and mock-infected cells. Five five-fold dilutions were made directly in the microtiter trays using a robot system. Untreated

-34-

virus controls, and HeLa cell controls were included in each test. Approximately 100 TCID₅₀ of Respiratory Syncytial Virus was added to two of the three rows in a volume of 50 µl. The same volume of medium was added to the third row to measure the cytotoxicity of the compounds at the same concentrations as those used to measure the antiviral activity. After two hours of incubation, a suspension (4×10^5 cells/ml) of HeLa cells was added to all wells in a volume of 50µl. The cultures were incubated at 37°C in a 5% CO₂ atmosphere. Seven days after infection the cytotoxicity and the antiviral activity was examined spectrophotometrically. To each well of the microtiter tray, 25 µl of a solution of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) was added . The trays were further incubated at 37°C for 2 hours, after which the medium was removed from each cup. Solubilization of the formazan crystals was achieved by adding 100 µl 2-propanol. Complete dissolution of the formazan crystals were obtained after the trays have been placed on a plate shaker for 10 min. Finally, the absorbances were read in an eight-channel computer-controlled photometer (Multiskan MCC, Flow Laboratories) at two wavelengths (540 and 690 nm). The absorbance measured at 690 nm was automatically subtracted from the absorbance at 540 nm, so as to eliminate the effects of non-specific absorption.